

Flora of the Pacific Northwest Checklist

A checklist of the Vascular Plants of the Pacific Northwest
Hosted by the University of Washington Herbarium

Family: Crassulaceae

24 terminal taxa (species, subspecies, and varieties).

The Flora of the Pacific Northwest Checklist aims to be a complete list of the native and naturalized vascular plants of the Pacific Northwest, with current classifications, nomenclature and synonymy.

Taxa included in the checklist:

- Native taxa whether extant, extirpated, or extinct.
- Exotic taxa that are naturalized, escaped from cultivation, or persisting wild.
- Waifs (e.g., ballast plants, escaped crop plants) and other scarcely collected exotics.
- Interspecific hybrids that are frequent or self-maintaining.
- Some unnamed taxa in the process of being described.

Family classifications follow [APG IV](#) for angiosperms, PPG I (J. Syst. Evol. 54:563-603. 2016.) for pteridophytes, and Christenhusz et al. (Phytotaxa 19:55-70. 2011.) for gymnosperms, with a few exceptions. Nomenclature and synonymy at the rank of genus and below follows the [2nd Edition of the Flora of the Pacific Northwest](#) except where superseded by new information.

Accepted names are indicated with **blue type**, synonyms with **gray type**.

Native species and infraspecies are marked with **bold-face type**.

*Non-native and introduced taxa are preceded by an asterisk.

Please note: This is a working checklist, continuously updated. Use it at your discretion.

Created from the Flora of the PNW Checklist database on April 4th, 2026 at 1:16am PT.

Available online at <https://burkeherbarium.org/pnwflora/>

Comments and questions should be addressed to the checklist administrators:

David Giblin (dgiblin@uw.edu)

Peter Zika (zikap941@gmail.com)

Suggested citation:

LASTNAME, F., OTHER PEOPLE. 2023+. Checklist of the Vascular Plants of the Pacific Northwest. University of Washington Herbarium. <https://www.pnwherbaria.org/pnwflora/>. Accessed Apr 4, 2026.

Dicots:

Crassulaceae [FNA8, HC, HC2] Stonecrop Family

Crassula [FNA8, HC2]

Sp. Pl. 1: 282. 1753; Gen. Pl. ed. 5, 136. 1754.
pygmy-weed

Tillaea [HC]

Crassula aquatica (L.) Schönland [FNA8, HC2]

Nat. Pflanzenfam. 51[III,2a]: 37. 1890.
wrinkle-seed pygmyweed, water pygmy weed

Crassula saginoides (Maxim.) M. Bywater & Wickens [KZ99]

Crassula vaillantii (Willd.) Roth., misapplied

Hydrophila vaillantii House

Tillaea angustifolia Nutt. var. *bolanderi* S. Watson

Tillaea aquatica L. [HC]

Tillaea bolanderi (S. Watson) Greene

Tillaea drummondii Torr. & A. Gray var. *bolanderi* (S. Watson) Jeps.

Tillaea vaillantii A. Gray, invalid name

Tillaeastrum pringlei Rose

FNA8: "The typical form of *Crassula aquatica*, with very short fruiting pedicels, grows chiefly in coastal salt marsh. It is rare and widely scattered in Alaska and Canada (W. J. Cody 1954), scarcely more common southward. M. Bywater and G. E. Wickens (1984) separated *C. saginoides* by pedicels elongate in fruit, sometimes to ca. 2 cm. It grows mostly inland and sometimes to 3000 meters, but from the specimens that they annotated, the ranges are not distinct. N. L. Britton and J. N. Rose (1905) and W. L. Jepson (1923-1925) have separated it, at least varietally, under other names, but most authors have included it without comment or at most have called it doubtfully distinct. I call it merely a phase of *C. aquatica* not needing a formal name (R. V. Moran 1992b). A typical strand plant is depicted in the lower left corner of the illustration panel on this page."

Crassula connata (Ruiz & Pav.) A. Berger [FNA8, HC2]

Nat. Pflanzenfam. ed. 2. 18a: 389. 1930.
pygmy weed

Crassula connata (Ruiz & Pav.) A. Berger var. *subsimplex* (S. Watson) M. Bywater & Wickens [JPM2]

Crassula erecta (Hook. & Arn.) A. Berger

Tillaea connata Ruiz & Pav.

Tillaea erecta Hook. & Arn.

Tillaea leptosepala Benth.

FNA8: "M. Bywater and G. E. Wickens (1984) proposed five varieties, four partly in the flora area, where they have largely overlapping ranges and overlapping characteristics. Further, some supposed differences probably result from local and year-to-year rainfall differences. Although probably having some genetic basis, these varieties seem too poorly defined to be useful (R. V. Moran 1992b). S. L. Hatch et al. (1990) were first to report *Crassula connata* in Texas, from a 1968 collection (F. B. Jones 7292), suggesting that it may be a recent arrival there. It was first discovered in British Columbia and Washington in 1977 (A. Ceska and O. Ceska 1980) and is treated as a rare native species by the heritage program in British Columbia and the Washington Natural Heritage Program, where it has a habitat typical of other southern disjunct natives (A. Ceska, pers. comm.)."

Crassula solieri (Gay) F. Meigen [FNA8, HC2]

Bot. Jahrb. Syst. 17: 239. 1893.
smooth-seed pygmyweed
(see also *Crassula aquatica*)

Tillaea solieri Gay

Recently (2017) collected along Columbia River in central WA.

**Crassula tillaea* Lester-Garl. [FNA8, HC2]

Fl. Jersey. 87. 1903.
mossy stonecrop

Tillaea muscosa L.

FNA8: "First found in California in 1925 (J. T. Howell 1942), *Crassula tillaea* is now widespread and locally common, often mingling with *C. connata*. It has been spreading northward, being first found in Oregon in 1984 (D. H. Wagner 1991), in Washington in 1999 (A. L. Jacobson et al. 2001), and in British Columbia in 2002 (P. F. Zika 2002); it has not yet been found growing with *C. connata* in this part of its range (A. Ceska, pers. comm.)."

Crassula viridis (S. Watson) Fedde [FNA8, HC2]

Just's Bot. Jahresber. 31(1): 829. 1904.
pricklyseed pygmy-weed

Rhodiola [FNA8, HC2]

Sp. Pl. 2: 1035. 1753; Gen. Pl. ed. 5, 457. 1754.
roseroot

Rhodiola integrifolia Raf. [FNA8, HC2]

Atlantic J. 1: 146. 1832.
king's crown, midsummer-men, roseroot

Sedum integrifolium (Raf.) A. Nelson

ssp. integrifolia [FNA8, HC2]

Atlantic J. 1(4): 146. 1832.
king's crown, roseroot

Sedum alaskanum (Rose) Rose ex Hutch.

Sedum roseum (L.) Scop. *ssp. integrifolium* (Raf.) Hultén [JPM], orthographic variant

FNA8: "The plants treated here as *Rhodiola integrifolia* and *R. rosea* are part of a difficult polymorphic complex of arctic to cool-temperate North America and Eurasia and of high mountains southward. Some authors have included them all in *R. rosea* [or *Sedum rosea* (Linnaeus) Scopoli], often with subspecies or varieties; N. L. Britton and J. N. Rose (1905) earlier divided them into two to several species. For this complex C. H. Uhl (1952) cited six published chromosome counts from Greenland through Eurasia to Japan, all $n = 11$ or $2n = 22$; he found the same numbers in seven collections from northeastern North America (all these *Rhodiola rosea* proper). From Eurasia, according to R. L. Taylor and G. A. Mulligan (1968), races with $2n = 16$ and 33 also are known. On the other hand, for endemics in Minnesota and New York and for five plants from New Mexico and California, Uhl found $n = 18$ or $2n = 36$, and Taylor and Mulligan likewise found $2n = 36$ in plants of Moresby Island, British Columbia. With the support of five more counts, but with none for the large area of Oregon and Wyoming to the Bering Sea, R. T. Clausen (1975) separated the 36-chromosome plants as *Sedum integrifolium*. More counts of $2n = 36$ have since appeared, including one from Sutwick Island, off the Alaska Peninsula (Å. Löve 1979). In middle North America, *Rhodiola integrifolia* and *R. rosea* are geographically distinct. The local endemic subsp. *leedyi* of the former grows in Minnesota, midway between the western subspecies of *R. integrifolia* and the eastern *R. rosea*, and grows in New York state within 100 km of *R. rosea*. Otherwise, the ranges of the two species are over 2000 km apart in the south and nearly 3000 km in the north. *Rhodiola integrifolia* also is the prevailing plant in eastern Asia, where it has been named *Sedum atropurpureum* N. S. Turczaninow (E. Hultén 1941-1950, vol. 5), and *R. rosea* seems to extend (although not verified by chromosome counts) from eastern Asia to far-western Alaska, on the coast of the Bering Sea. Although saying that *Sedum integrifolium* differs from *S. rosea* in many ways besides the chromosome number, R. T. Clausen (1975) found few absolute distinctions. His best key characters were those used here, petal width of staminate flowers, largely supported by flower color. Although questions remain unanswered, it seems best for now to follow Clausen in keeping the two species for North America. Over its broad range, *Rhodiola integrifolia* is quite variable (e.g., see E. Hultén 1941-1950, vol. 5). R. T. Clausen (1975) noted that in some populations pistillate plants outnumber staminate; in others staminate may be six times as many as pistillate. He distinguished two outlying endemics as subspp. *leedyi* and *neomexicana*, also kept as subspecies

here. He also proposed subsp. *procer[a]* for tall robust plants of Colorado, New Mexico, and (less typical) California, all within the range of subsp. *integrifolia* and all with the same chromosome number. Some of his plants look remarkably different from the usual dwarf forms of subsp. *integrifolia* that grow at the same high elevations. He did not include in subsp. *procera* (and apparently did not see alive) the tall plants often found inland in Alaska and northwestern Canada, which would be *Sedum frigidum* Rydberg according to Hultén. Thus the racial situation is much more complex than the naming of only two peripheral subspecies might suggest."

Rhodiola rhodantha (A. Gray) H. Jacobsen [FNA8, HC2]

Natl. Cact. Succ. J. 28: 5. 1973.

rose crown

Sedum rhodanthum A. Gray [HC]

Sedum [FNA8, HC, HC2]

Sp. Pl. 1: 430. 1753; Gen. Pl. ed. 5, 197. 1754.

stonecrop

(see also *Rhodiola*)

****Sedum acre*** L. [FNA8, HC, HC2]

Sp. Pl. 1: 432. 1753.

mossy stonecrop

Sedum elrodii M.E. Jones

****Sedum album*** L. [FNA8, HC2]

Sp. Pl. 1: 432. 1753.

white stonecrop

FNA8: "Sedum album was first reported as naturalized in the United States in 1934."

Sedum borschii (R.T. Clausen) R.T. Clausen [FNA8, HC2]

Sedum N. Amer. 298. 1975.

Borsch's stonecrop

****Sedum brevifolium*** DC. [HC2]

Soc. Agr. Dept. Seine 11: 79. 1808.

short-leaved stonecrop

Recently collected in King County, where fully naturalized on a montane rocky bald north of Interstate 90 east of North Bend. The bald is adjacent to rock climbing routes, suggesting propagules may have arrived with climbers. The plants form reproducing populations across a several hectare area of balds and rock faces.

Sedum debile S. Watson [FNA8, HC, HC2]

Botany (Fortieth Parallel). 102. 1871.

weakstemmed stonecrop

Sedum divergens S. Watson [FNA8, HC, HC2]

Proc. Amer. Acad. Arts. 17: 372. 1882.

Pacific stonecrop, spreading stonecrop

Amerosedum divergens (S. Watson) Á. Löve & D. Löve

FNA8: "Leaves of *Sedum divergens* are close-set, thick, and turgid. This species occurs in scattered and disjunct populations from the coastal mountains of Alaska (D. F. Murray 1980) and British Columbia to the northern Cascade Mountains and Olympic Mountains of Washington; Lake Peak, Josephine County, Oregon; and Klamath Mountains near Mount Robson in Alberta and British Columbia."

****Sedum forsterianum*** Sm. [HC2]

Engl. Bot. 26: t. [1 Oct 1807-1 Apr 1808]. 1802.

Forster's stonecrop

Sedum lanceolatum Torr. [FNA8, HC, HC2]

Ann. Lyceum Nat. Hist. New York. 2: 205. 1827.

lance-leaved stonecrop

(see also *Sedum rupicola*)

Amerosedum nesioticum (G.N. Jones) Á. Löve & D. Löve
Sedum lanceolatum Torr. ssp. *lanceolatum* [KZ99]
Sedum lanceolatum Torr. ssp. *nesioticum* (G.N. Jones) R.T. Clausen [KZ99]
Sedum lanceolatum Torr. var. *lanceolatum* [FNA8, HC]
Sedum lanceolatum Torr. var. *nesioticum* (G.N. Jones) C.L. Hitchc. [FNA8, HC]
Sedum nesioticum G.N. Jones
Sedum stenopetalum Pursh var. *subalpinum* Fröd.

***Sedum leibergii* Britton [FNA8, HC, HC2]**

N. Amer. Fl. 22: 73. 1905.
Leiberg's stonecrop

Amerosedum leibergii (Britton) Á. Löve & D. Löve
Sedum divaricatum S. Watson

***Sedum oreganum* Nutt. [FNA8, HC, HC2]**

Fl. N. Amer. 1: 559. 1840.
Oregon stonecrop

Gormania oregana (Nutt.) Britton
Sedum oreganum Nutt. ssp. *oreganum* [KZ99]
Sedum oreganum Nutt. ssp. *tenue* R.T. Clausen
Sedum oreganum Nutt. var. *oreganum* [FNA8]
Sedum oreganum Nutt. var. *tenue* (R.T. Clausen) H. Ohba [FNA8]

***Sedum oregonense* (S. Watson) M. Peck [FNA8, HC, HC2]**

Man. Higher Pl. Oregon. 361. 1941.
creamy stonecrop

***Sedum rupicola* G.N. Jones [FNA8, HC2]**

Res. Stud. State Coll. Wash. 2: 125. 1931.
lance-leaved stonecrop

Sedum lanceolatum Torr. var. *rupicola* (G.N. Jones) C.L. Hitchc., orthographic variant
Sedum lanceolatum Torr. var. *rupiculum* (Jones) Hitchc. [HC]

FNA8: "The leaves of *Sedum rupicola* detach very easily and the fallen ones sprout and produce plantlets from their bases. R. T. Clausen (1975) considered *S. rupicola* to be most closely related to *S. lanceolatum*. He recognized it as a species because, although it sometimes grows sympatrically with *S. lanceolatum*, the two do not hybridize, and because *S. rupicola* flowers a week earlier and grows in soils of higher pH than does *S. lanceolatum*. The general morphological differences are: in *S. rupiculum* leaves of sterile shoots are ovate and detach easily, sepals have obtuse apices, petals have minutely mucronate tips (0.1 mm), and nectaries are deep yellow; in *S. lanceolatum* leaves of sterile shoots are linear-lanceolate and do not detach easily, sepals have acute apices, petal apices are long-acuminate (0.8 mm), and nectaries are pale yellow."

***Sedum spathulifolium* Hook. [FNA8, HC, HC2]**

Fl. Bor.-Amer. 1: 227. 1832.
broadleaf stonecrop, spatula-leaf stonecrop

Sedum pruinosum Britton
Sedum spathulifolium Hook. ssp. *pruinum* (Britton) R.T. Clausen & C.H. Uhl
Sedum spathulifolium Hook. ssp. *spathulifolium* [KZ99]
Sedum spathulifolium Hook. var. *pruinum* (Britton) B. Boivin [FNA8]
Sedum spathulifolium Hook. var. *spathulifolium* [FNA8]

***Sedum stenopetalum* Pursh [FNA8, HC, HC2]**

Fl. Amer. Sept. 1: 324. 1813.
wormleaf stonecrop

Amerosedum stenopetalum (Pursh) Á. Löve & D. Löve
Sedum douglasii Hook.

ssp. *ciliolum* (Howell) R.T. Clausen [HC2]

Sedum radiatum S. Watson var. *ciliolum* (Howell) H. Ohba [FNA8]

ssp. *stenopetalum* [HC2, KZ99]

wormleaf stonecrop

Sedum monanthum Suksd.

Sedum stenopetalum Pursh ssp. *monanthum* (Suksd.) R.T. Clausen [KZ99]

Sedum stenopetalum Pursh var. *monanthum* (Suksd.) H. Ohba [FNA8]

Sedum stenopetalum Pursh var. *stenopetalum* [FNA8]

****Sedum thartii*** L.P. Hébert [HC2]

Rev. Cytol. Biol. Vég. Bot. 6(3?4): 211 (1983), nom. nov., as 'stat. nov.': (1983). 1983.

Thart's stonecrop

Sedum rupestre L. [FNA8], misapplied

Sporadically naturalized on road cuts and other disturbed areas in western Washington and southwestern B.C. FNA8: "Most naturalized records of *S. rupestre* in North America have been incorrectly named *S. reflexum*." Gallo and Zika (2014) determined that the names *Sedum rupestre* and *S. reflexum* are misapplied to North American plants; our plants can be assigned to *Sedum thartii* and *S. forsterianum*.

Sedum valens Björk [HC2]

Idaho stonecrop