

Flora of the Pacific Northwest Checklist

A checklist of the Vascular Plants of the Pacific Northwest
Hosted by the University of Washington Herbarium

Family: Malvaceae

34 terminal taxa (species, subspecies, and varieties).

The Flora of the Pacific Northwest Checklist aims to be a complete list of the native and naturalized vascular plants of the Pacific Northwest, with current classifications, nomenclature and synonymy.

Taxa included in the checklist:

- Native taxa whether extant, extirpated, or extinct.
- Exotic taxa that are naturalized, escaped from cultivation, or persisting wild.
- Waifs (e.g., ballast plants, escaped crop plants) and other scarcely collected exotics.
- Interspecific hybrids that are frequent or self-maintaining.
- Some unnamed taxa in the process of being described.

Family classifications follow [APG IV](#) for angiosperms, PPG I (J. Syst. Evol. 54:563-603. 2016.) for pteridophytes, and Christenhusz et al. (Phytotaxa 19:55-70. 2011.) for gymnosperms, with a few exceptions. Nomenclature and synonymy at the rank of genus and below follows the [2nd Edition of the Flora of the Pacific Northwest](#) except where superseded by new information.

Accepted names are indicated with **blue type**, synonyms with **gray type**.

Native species and infraspecies are marked with **bold-face type**.

*Non-native and introduced taxa are preceded by an asterisk.

Please note: This is a working checklist, continuously updated. Use it at your discretion.

Created from the Flora of the PNW Checklist database on April 4th, 2026 at 1:14am PT.

Available online at <https://burkeherbarium.org/pnwflora/>

Comments and questions should be addressed to the checklist administrators:

David Giblin (dgiblin@uw.edu)

Peter Zika (zikap941@gmail.com)

Suggested citation:

LASTNAME, F., OTHER PEOPLE. 2023+. Checklist of the Vascular Plants of the Pacific Northwest. University of Washington Herbarium. <https://www.pnwherbaria.org/pnwflora/>. Accessed Apr 4, 2026.

Dicots:

Malvaceae [HC, HC2] Mallow Family

**Abutilon* [HC2]

Indian mallow

**Abutilon theophrasti* Medik. [HC2, JPM]

Malvenfam. 28. 1787.

velvetleaf

Noxious.

**Alcea* [HC2]

hollyhock

**Alcea rosea* L. [HC2, JPM2]

Sp. Pl. 2: 687. 1753.

hollyhock

Althaea rosea (L.) Cav.

The specimens for this species in Washington are from cultivated settings. It is not considered naturalized in the flora.

**Gossypium* [HC2]

**Gossypium hirsutum* L. [HC2]

cotton

Collected twice in Washington along roadsides. No evidence that self-sustaining populations exist in Washington.

**Hibiscus* [HC, HC2]

rosemallow

**Hibiscus moscheutos* L. [HC2, JPM]

Sp. Pl. 2: 693. 1753.

crimson-eyed rosemallow

This species is known from a single location in Washington as a garden escape. More collections are needed to demonstrate that it is established in the flora.

**Hibiscus trionum* L. [HC, HC2]

Sp. Pl. 2: 697. 1753.

flower-of-an-hour

Trionum trionum (L.) Wooton & Standl.

All specimens for this species in Washington are from cultivated settings. This species is considered excluded until specimens are located that indicate that it is established in the flora in Washington.

Iliamna [HC, HC2]

globemallow

Iliamna longisepala (Torr.) Wiggins [HC, HC2]

Leafl. Bot. Observ. Crit. 1(15): 206. 1906.

long-sepal globemallow

Iliamna rivularis (Douglas) Greene [HC, HC2]

Leafl. Bot. Observ. Crit. 1: 206. 1906.

streambank globemallow

Iliamna rivularis (Douglas) Greene var. *diversa* (A. Nelson) Wiggins [HC]

Iliamna rivularis (Douglas) Greene var. *rivularis* [HC]
Sphaeralcea rivularis (Douglas) Torr.

**Malva* [HC, HC2]

cheeses, cheeseweed, mallow

**Malva alcea* L. [HC2]

**Malva moschata* L. [HC, HC2]

Sp. Pl. 2: 690. 1753.

musk mallow

**Malva neglecta* Wallr. [HC, HC2]

Sylloge Plantarum Novarum 1: 140-142. 1824.

dwarf mallow

Malva rotundifolia L., misapplied

**Malva parviflora* L. [HC, HC2]

Demonstr. Pl. 18. 1753.

cheeseweed, alkali mallow, small-whorl mallow

**Malva pusilla* Sm. [HC2]

English Botany 4: pl. 241. 1795.

low mallow

**Malva sylvestris* L. [HC, HC2]

Sp. Pl. 2: 689. 1753.

common mallow, high mallow

**Malva verticillata* L. [HC2]

This species is known from Washington by a single collection in 1946. It is considered excluded until additional specimens indicate that it is established in the flora.

**Malvastrum* [HC2]

**Malvastrum coromandelianum* (L.) Garcke [HC2]

Malvella [HC2]

alkali-mallow

Malvella leprosa (Ortega) Krapov. [HC2, JPM]

Bonplandia (Corrientes) 3(5): 59. 1970.

alkali-mallow

Sida hederacea (Douglas ex Hook.) Torr. ex A. Gray [HC]

Sida leprosa (Ortega) K. Schum. var. *hederacea* (Douglas) K. Schum. ex Clement

**Modiola* [HC2]

**Modiola caroliniana* (L.) G. Don [HC2]

**Sida* [HC, HC2]

**Sida spinosa* L. [HC2]

Sidalcea [HC, HC2]

checker-mallow, sidalcea

Sidalcea asprella Greene [HC2]

ssp. *asprella* [HC2]

**Sidalcea campestris* Greene [HC, HC2]

Bulletin of the California Academy of Sciences 1(3): 76-77. 1885.

meadow checker-mallow, meadow sidalcea

Sidalcea asplenifolia Greene
Sidalcea sylvestris A. Nelson

Native only to the Willamette Valley area (Multnomah and Washington to Benton and Linn counties). WA specimens were collected by Piper in late 1800s near Seattle. These specimens are considered likely introductions. Piper and Beattie's 1915 "Flora of the Northwest Coast contains this note for *S. campestris*: "In moist meadows, Willamette Valley, Oregon. *S. asplenifolia* Greene found at Seattle in hay meadows is apparently the same and perhaps was introduced with grass seed." The lack of specimens for this species from WA over the last 100 years strongly suggests that it is likely not part of the contemporary flora. It is best considered an occasional garden escape.

Sidalcea cusickii Piper [HC, HC2]

Cusick's checker-mallow

Sidalcea hendersonii S. Watson [HC, HC2]

Proceedings of the American Academy of Arts and Sciences 23(2): 262. 1888.

Henderson's checker-mallow

Sidalcea hirtipes C.L. Hitchc. [HC, HC2]

Perenn. Sp. *Sidalcea* 42-44, map 2. 1957.

bristly-stem checker-mallow

Rare in Washington.

Sidalcea nelsoniana Piper [HC, HC2]

Proc. Biol. Soc. Wash. 32(12): 41. 1919.

Nelson's checker-mallow

Known from Cowlitz and Lewis counties.

Sidalcea neomexicana A. Gray [HC, HC2]

Rocky Mountain checker-mallow

Sidalcea neomexicana A. Gray var. *crenulata* (A. Nelson) C.L. Hitchc. [HC]

Sidalcea oregana (Nutt. ex Torr. & A. Gray) A. Gray [HC, HC2]

Oregon checker-mallow

var. *calva* C.L. Hitchc. [HC, HC2]

Perenn. Sp. *Sidalcea* 61, map 3. 1957.

Oregon checkermallow

Listed as rare by WNHP.

var. *oregana* [FNA12, HC, HC2]

Memoirs of the American Academy of Arts and Science, new series 4(1): 20. 1849.

Oregon checkermallow

Sidalcea oregana (Nutt. ex Torr. & A. Gray) A. Gray ssp. *oregana*

Sidalcea oregana (Nutt. ex Torr. & A. Gray) A. Gray var. *maxima* (M. Peck) C.L. Hitchc. [HC]

Sidalcea oregana (Nutt. ex Torr. & A. Gray) A. Gray var. *procera* C.L. Hitchc. [HC]

FNA12: "Subspecies *oregana* is variable; it intergrades with subsp. *spicata* and *Sidalcea setosa*. C. L. Hitchcock (1957) accepted five varieties within the typical subspecies; morphological intergrades exist. A case can be made for recognition of var. *calva* C. L. Hitchcock, which has been listed as endangered both federally and in Washington, where it is endemic. These plants are generally robust, sparsely appressed-hairy with four-rayed hairs, the leaves are generally nearly glabrous and fleshy-textured, and the calyx lobes are subglabrous and ciliate. Found in the Wenatchee Mountains, an area of high endemism, var. *calva* does not appear to be much different from other, nearly glabrous populations elsewhere; it appears to be the only variety with a chromosome count of $2n = 60$. This treatment does not accept both subspecies and varieties within *Sidalcea*; therefore, it has been placed here into synonymy with the wide-ranging, variable typical subspecies. Subspecies *oregana* can generally be distinguished from the other subspecies by its more-open inflorescences that are elongated in fruit, its multistemmed clumps as much as 30 cm in diameter, its generally stellate-hairy to glabrescent stem bases, its generally uniformly stellate-hairy calyces, its somewhat reticulate-roughened mericarps, and its bracts that are generally equal to or shorter than the young flower buds. Subspecies *oregana* appears to be the source of commonly sold cultivars. It has been

listed as sensitive in Montana and as rare in British Columbia."

var. *spicata* (Regel) Jeps. [HC, HC2]

Sidalcea virgata Howell [HC, HC2]

Fl. N.W. Amer. 1: 101. 1897.

virgate checker-mallow, virgate checkerbloom

Sidalcea malviflora (DC.) A. Gray ex Benth. ssp. *virgata* (Howell) C.L. Hitchc. [KZ99]

Sidalcea malviflora (DC.) A. Gray ex Benth. var. *virgata* (Howell) Dimling

Extirpated from Washington according to WNHP. One specimen (Ed Alverson, 1987) at WTU from a population in Thurston County. Otherwise only known from Oregon.

Sphaeralcea [HC, HC2]

globe-mallow

Sphaeralcea coccinea (Nutt.) Rydb. [HC, HC2]

red globe-mallow

var. *coccinea* [HC2]

Sphaeralcea grossulariifolia (Hook. & Arn.) Rydb. [HC, HC2]

Bull. Torrey Bot. Club 40(2): 58. 1913.

gooseberry-leaved globemallow

Sphaeralcea grossulariaefolia (Hook. & Arn.) Rydb. [HC], orthographic variant

Sphaeralcea grossulariifolia (Hook. & Arn.) Rydb. ssp. *grossulariifolia* [KZ99]

Sphaeralcea grossulariifolia (Hook. & Arn.) Rydb. var. *moorei* S.L. Welsh

Sphaeralcea munroana (Douglas ex Lindl.) Spach [HC, HC2]

Proc. Amer. Acad. Arts 22(2): 292. 1887.

Munro's globemallow, white-stemmed globemallow

Sphaeralcea munroana (Douglas ex Lindl.) Spach ssp. *munroana* [JPM]

****Tilia*** [HC2]

small-leaved linden

****Tilia cordata*** Mill. [HC2]

Gard. Dict., ed. 8. n. 1. 1768.

small-leaved linden

****Tilia platyphyllos*** Scop. [HC2]

All specimens for this species in Washington are from cultivated settings. It is not considered established in the flora.